



This Product Information Sheet is not intended to replace the Product Label. Always read the Product Label for the most current nutrient content and feeding directions.

14:1:14 C2800 ROL.433 Mineral

Product number: 10918



Product Information

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

For control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline in beef cattle, for control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella spp.* susceptible to chlortetracycline in beef cattle and prevention of the development of face flies, horn flies, house flies, and stable flies in the manure of treated cattle.

Feeding Instructions

WARNING: Withdraw 48 hours prior to slaughter. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

CAUTION: Use as directed. Consumption of selenium should not exceed 3 mg per head per day.

WARNING: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

Provide fresh, clean water to all animals at all times.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

USER SAFETY RECOMMENDATIONS

Personal Protective Equipment (PPE): Long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS: This compound is toxic to fish.

Storage and Disposal: Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a dry place in original container.

Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Feeding and Management

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For control of bacterial pneumonia and control of flies in cattle up to 700 lbs.: Thoroughly mix 500 lbs. SWEETLIX 14:1:14 High Mag with CTC and Rabon Oral Larvacide with 1,500 lbs. of non-medicated feed. Feed one pound of this medicated mixture to supply 350 mg CTC and 490 mg Rabon per head per day. Cattle should receive 350 mg CTC and 70 mg Rabon per 100 lbs. of body weight per day (2.1 grams per 100 lbs. body weight per month) for the stated drug claims.

For control of anaplasmosis and control of flies in beef cattle up to 700 lbs.: Thoroughly mix 500 lbs. SWEETLIX 14:1:14 High Mag with CTC

and Rabon Oral Larvacide with 1,500 lbs. of non-medicated feed. Feed one pound of this medicated mixture to supply 350 mg CTC and 490 mg Rabon per head per day. Cattle should receive 350 mg CTC and 70 mg Rabon per 100 lbs. of body weight per day (2.1 grams per 100 lbs. body weight per month) for the stated drug claims.

For control of anaplasmosis and control of flies in beef cattle up to 1,000 lbs.: Thoroughly mix 714 lbs. SWEETLIX 14:1:14 High Mag with CTC and Rabon Oral Larvacide with 1,286 lbs. of non-medicated feed. Feed one pound of this medicated mixture to supply 500 mg CTC and 700 mg Rabon per head per day. Cattle should receive 500 mg CTC and 70 mg Rabon per 100 lbs. of body weight per day (2.1 grams per 100 lbs. body weight per month) for the stated drug claims.

Fly Control Program

SWEETLIX 14:1:14 High Mag with CTC and Rabon Oral Larvacide prevents the development of horn flies, house flies and stable flies in the manure of treated cattle, but is not effective against existing adult flies. When starting a feeding program during the fly season it may be necessary to use other control measures to reduce the population of existing adult flies. Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which breed not only in other decaying matter or silage on the premises. Start feeding SWEETLIX 14:1:14 High Mag with CTC and Rabon Oral Larvacide early in the spring before flies begin to appear and continue feeding throughout the summer and into fall. SWEETLIX 14:1:14 High Mag with CTC and Rabon Oral Larvacide may be fed to cattle up to slaughter.

Rabon® is a registered trademark of Bayer HealthCare, LLC.

Nutrition Information

Calcium, Min	14.00%
Calcium, Max	16.80%
Phosphorus, Min	1.00%
Salt, Min	19.00%
Salt, Max	22.80%
Magnesium, Min	14.00%
Copper, Min	200 ppm
Iodine, Min	5 ppm
Manganese, Min	850 ppm
Selenium, Min	26 ppm
Zinc, Min	575 ppm
Vitamin A, Min	100,000 IU/lb
Vitamin D-3, Min	10,000 IU/lb
Vitamin E, Min	10 IU/lb

Product Ingredients

Active Drug Ingredients:

Chlortetracycline.....2,800 g/ton

Tetrachlorvinphos.....0.433%

Calcium Carbonate, Salt, Magnesium Oxide, Molasses Products, Monocalcium Phosphate, Dicalcium Phosphate, Roughage Products, Processed Grain By-Products, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Red Iron Oxide and Artificial Flavoring.