



This Product Information Sheet is not intended to replace the Product Label. Always read the Product Label for the most current nutrient content and feeding directions.

## 14:1:14 ROL.494 Mineral

Product number: 10925



### Product Information

For control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies, and stable flies in the manure of treated cattle

- Self-fed supplement for beef cattle on pasture or with all-roughage rations
- Scientifically balanced to be fed free-choice to cattle
- All protein, mineral and vitamin ingredients are selected for high biological availability and stability

*Rabon® is a registered trademark of Bayer HealthCare, LLC.*

### Feeding Instructions

#### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals:**

#### CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

**Environmental Hazards:** This compound is toxic to fish.

#### USER SAFETY RECOMMENDATIONS

**Personal Protective Equipment (PPE):** Long-sleeved shirt and long pants, shoes and socks, chemical resistant gloves. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Follow manufacturers instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Storage and Disposal:** Do not contaminate water, food, or feed by storage or disposal.

**Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and Local Authorities, by burning. If burned, stay out of smoke. Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**Follow feeding directions.** Consumption of selenium should not exceed 3 mg per head daily.

#### Directions for Use

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling.*

**USE RESTRICTIONS:** Not to be used on horses destined for slaughter.

**PRECAUTIONS FOR HORSE USE:** This product contains tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, consult your veterinarian immediately. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.

#### GENERAL INSTRUCTIONS

SWEETLIX 14:1:14 High Mag with Rabon should be fed continuously on a free choice basis to cattle to provide minerals, vitamins and Rabon daily. Feed as the only FREE CHOICE source of salt, other minerals or vitamins. Rabon Oral Larvacide should be fed at a level to provide 70 mg Rabon per 100 lb. of body weight per day (2.1 grams per 100 lbs. body weight per month). Feed supplies (pasture or other forages) and feeder locations should be adequate to insure proper mineral consumption. Consumption in accordance with intake levels listed below will deliver 70 mg Rabon per 100 lb. body weight per day.

Animal

Daily Intake

Weight (lb)	Supplement Intake (oz)	(lb)	of Rabon® (mg)
500	2.5	0.15625	350
600	3.0	0.18750	420
700	3.5	0.21875	490
800	4.0	0.25000	560
900	4.5	0.28125	630
1000	5.0	0.31250	700
1100	5.5	0.34375	770
1200	6.0	0.37500	840

If animals consume more or less than these levels, feeders should be moved further from or closer to the general resting or watering areas. If animals still fail to consume the correct amounts, discontinue feeding this free choice supplement.

Provide fresh, clean water to all animals at all times.

#### Fly Control Program

SWEETLIX 14:1:14 High Mag with Rabon prevents the development of fecal flies in the manure of treated cattle, but it is not effective against existing adult flies. It may be necessary to use other control measures to reduce the population of existing adult flies.

Supplemental fly control measures may be needed in and around buildings to control adult house flies and stable flies which breed not only in animal manure but also in other decaying matter on the premises. In order to achieve optimum fly control, SWEETLIX 14:1:14 High Mag with Rabon should be used in conjunction with other good management and sanitation practices. Start feeding

SWEETLIX 14:1:14 High Mag with Rabon early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall. SWEETLIX 14:1:14 High Mag with Rabon may be fed to cattle up to slaughter.

## Nutrition Information

Calcium, Min	13.00%
Calcium, Max	15.00%
Phosphorus, Min	1.00%
Salt, Min	24.00%
Salt, Max	26.00%
Magnesium, Min	14.00%
Copper, Min	200 ppm
Manganese, Min	860 ppm
Selenium, Min	16 ppm
Zinc, Min	570 ppm
Vitamin A, Min	20,000 IU/lb
Vitamin D-3, Min	2,000 IU/lb
Vitamin E, Min	2 IU/lb

## Product Ingredients

#### Active Drug Ingredient:

Tetrachlorvinphos.....0.494%

Other Ingredients.....99.506%

Calcium Carbonate, Magnesium Oxide, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Processed Grain By-Products, Molasses Products, Roughage Products, Manganous Oxide, Manganese Sulfate, Zinc Oxide, Zinc Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil, Artificial Flavoring and Red Iron Oxide.